

dna oestrogen

optimal health for life

Welcome

SUMMER

to your dna oestrogen report

Date of Birth: 30 Mar

Date Reported: 12 Jul 2023

Sample Number: M23001760

Referring Practitioner:

Welcome to your dna oestrogen report

From your DNA sample we have used a process called the Polymerase Chain Reaction (PCR), which copies the DNA of your genes many times over so that we can generate sufficient quantities to analyse your genetic material. We then identify unique DNA sequences in some of your genes.

Considerable inter-individual variability has been observed in biological areas that are involved in carcinogen metabolism, metabolism of steroid hormones, and phase I and phase II detoxification. Variations in genes involved in these biological processes help identify a sub-population of women and men with higher lifetime exposure to oestrogens, oestrogen metabolites and other carcinogens. Understanding an individual's genetic variability will allow for targeted diet, lifestyle and hormone intervention.

Understanding genetics

Before reading your full assessment, please take a few minutes to review this background information. This will help you better understand your results and enhance the value of this personalised report.

What are genes?

Genes are segments of DNA that contain the instructions your body needs to make each of the many thousands of proteins required for life. Each gene is comprised of thousands of combinations of "letters" which make up your genetic code. The code gives the instructions to make the proteins required for proper development and function.

What are gene variations?

With the exception of identical twins, all people have small differences (variations) in their genetic code. It is these differences that make each of us unique. An example of a genetic variation is that one "letter" may be replaced by another. These variations can lead to changes in the resulting proteins being made. For example a "C" may be changed to a "G" at a point in the genetic code. When the variation affects only one genetic "letter" it is called a Single Nucleotide Polymorphism, or SNP (pronounced "snip"). Variations can however also affect more than one "letter".

Are gene variations "bad"?

In general, variations should not be considered good or bad. Rather, genetic variations are simply slight differences in the genetic code. The key is to know which form of the variation you carry in order to make appropriate lifestyle choices.

How to read your results

You will find your genetic results in the following pages. On the left side you will see the gene name and description. On the right side you will find your specific result and an explanation of the results, associated risks, and diet and lifestyle recommendations. The impact can be identified by the colour of the circle (please see the key below).



No impact



Low impact



Moderate impact




High impact

Summary of results

Gene Name	Genetic Variation	Your Result	Gene Impact
CYP1A1	Msp1 T>C	TT	
CYP1A1	A>G Ile462Val	AA	
CYP1B1	C>G Val432Leu	CC	
CYP17A	34 T>C	TT	
MnSOD/SOD2	47 T>C (Val16Ala)	TT	
GSTM1	Insertion/Deletion	Present	
GSTT1	Insertion/Deletion	Present	
COMT	472 G>A (Val158Met)	AG	
MTHFR	677 C>T	CT	
SULT1A1	638 G>A Arg213His	GA	
NQ01	609 C>T	CC	
FACTOR V	G1691A	GG	

The combination of gene variants identified in this analysis indicates you have efficient oestrogen detoxification and no additional support is required.

Oestrogen metabolism & detoxification pathway



	Neutral /non-harmful metabolites
	Harmful metabolites
	Neutral/possibly harmful metabolites
	SNPs tested in DNA Oestrogen

Test results

CYP1A1 Msp1 T>C

The CYP1A1 gene encodes a phase I cytochrome P450 enzyme that converts environmental procarcinogens such as PAHs and aromatic amines to reactive intermediates having carcinogenic effects. In addition, CYP1A1 is involved in the oxidative metabolism of oestrogens, which may play a critical role in the aetiology of breast and prostate cancer. CYP1A1 enzyme catalyses the 2-hydroxylation of oestradiol (E1 and E2) in several extra hepatic tissues including breast tissue. It is also involved in activating cigarette smoke, diet and environmental pollutants, and producing carcinogens.

YOUR RESULT: TT

No variant was detected at the Msp1 T>C locus.

CYP1A1 Ile462Val A>G

The CYP1A1 gene encodes a phase I cytochrome P-450 enzyme that converts environmental procarcinogens such as PAHs and aromatic amines to reactive intermediates having carcinogenic effects. In addition, CYP1A1 is involved in the oxidative metabolism of oestrogens, which may play a critical role in the aetiology of breast and prostate cancer.

YOUR RESULT: AA

No variant was detected.

CYP1B1 1294 C>G

CYP1B1 enzymes catalyses the 4-hydroxylation of oestradiol, it also activates many PAHs and arylamines.

YOUR RESULT: CC

No variant was detected at the 1294 C>G locus.

CYP17A 34T>C

CYP17 mediates both steroid 17 α -hydroxylase and 17,20-lyase activities, and catalyses a rate-limiting step in ovarian and adrenal biosynthesis leading to the precursor, dehydroepiandrosterone. The C allele increases enzyme activity, thereby increasing the amount of bioavailable oestrogen.

YOUR RESULT: TT

No variant was detected at the 34 T>C locus.

NQO1 609 C>T

NADP(H:) quinone oxidoreductase 1 (NQO1) often referred to as Quinone Reductase is primarily involved in the detoxification of potentially mutagenic and carcinogenic quinones derived from tobacco smoke, diet and oestrogen metabolism. NQO1 also protects cells from oxidative stress by maintaining the antioxidant forms of ubiquinone and vitamin E.

YOUR RESULT: CC

The analysis identified no genetic variation at the 209 C>T locus.

GSTM1 Insertion/Deletion

Glutathione S-transferase M1 is the most biologically active member of the GST super-family and is involved in Phase II detoxification in the liver. It is responsible for the removal of xenobiotics, carcinogens, and products of oxidative stress. These enzymes are involved in the phase 2 conjugation of oestrogen quinones to glutathione.

YOUR RESULT: Present

The GSTM1 gene is present.

GSTT1 Insertion/Deletion

Glutathione S-transferases (GSTs) are a family of multifunctional enzymes involved in the metabolism of a variety of xenobiotic compounds, including mammary carcinogens. These enzymes are involved in the conjugation of oestrogen quinones to glutathione.

YOUR RESULT: Present

The GSTT1 gene is present.

COMT 472 G>A or Val158Met

Soluble catechol-O-methyltransferase (S-COMT) helps control the levels of certain hormones and is involved in methylation and inactivation of catechol oestrogens. Accumulation of oestrogen metabolites appears to confer increased risk of breast cancer via oxidative DNA damage.

YOUR RESULT: AG

The A allele is associated with a 3-4 fold reduction in the methylation activity of the COMT enzyme. For A allele carriers beneficial modulation of oestrogen metabolism can be accomplished through dietary and lifestyle modifications. Key interventions include increasing insoluble fibre, managing the quality of dietary fat intake, increasing phytoestrogen intake, losing weight, and increasing exercise. In addition, select nutrients and micronutrients effectively reduce oestrogen load by supporting preferred oestrogen pathways. These are included at the end of the report.

MTHFR 677 C>T

Methylenetetrahydrofolate Reductase (MTHFR) is a key enzyme in the folate metabolic pathway. Reduced activity influences the balance between DNA synthesis, repair and methylation processes.

YOUR RESULT: CT

The T allele lowers activity of the MTHFR enzyme, which results in an increase in homocysteine levels, a decrease in DNA methylation and an increase in DNA adducts. Enzyme function is 70% of optimal in CT genotype individuals. Decreased MTHFR enzyme activity has been associated with increased premenopausal breast cancer risk with long duration of oestrogen exposure. These individuals have increased folate, vitamin B2, B6 and B12 requirements. In addition to ensuring folate-rich foods, a general B vitamin or multi-vitamin supplement containing as much as 800ug folate may be recommended.

SULT1A1 638 G>A

Sulfotransferase 1A1 (SULT1A1) is involved in the inactivation of oestrogens and bio-activation of heterocyclic amines and polycyclic aromatic hydrocarbons.

YOUR RESULT: GA

A allele carriers have a substantially lower activity of this enzyme, and has been associated with a greater risk for post-menopausal breast cancer, increased with higher BMI, and longer exposure to endogenous hormones.

For individuals with the A allele beneficial modulation of oestrogen metabolism can be accomplished through dietary and lifestyle modifications. Key interventions include increasing insoluble fibre, avoiding refined CHO, increasing phytoestrogen intake, losing weight, and increasing exercise. In addition, select nutrients and micronutrients effectively reduce oestrogen load by supporting preferred oestrogen pathways. These are included at the end of the report.

MnSOD/SOD2 47T>C (Val16Ala)

The MnSOD/SOD2 enzyme destroys the free radicals which are normally produced within cells and which are damaging to biological systems. The enzyme thus has important anti-oxidant activity within the cell, especially within the mitochondria.

YOUR RESULT: TT

There is evidence that people without the variant, i.e. those without the T allele, and with a lower consumption of fruits and vegetables, are at increased risk of developing disease, including the risk of developing breast cancer.

FACTOR V G1691A

Factor V functions as a cofactor to allow factor Xa to activate the enzyme thrombin, and in turn cleaves fibrinogen to form fibrin, which polymerizes to form the dense meshwork that makes up the majority of a clot. Activated protein C (aPC) is a natural anticoagulant that acts to limit the extent of clotting by cleaving and degrading factor V. Factor V Leiden gene mutation is characterised by a poor anticoagulant response to APC and an increased risk for venous thromboembolism (VTE). Deep venous thrombosis (DVT) is the most common VTE, with the legs being the most common site however it VTE can also occur in other parts of the body including the brain, eyes, liver, and kidneys.

YOUR RESULT: GG

No variant was detected at the 1691 G>A locus.

Nutrition and oestrogen

If a moderate or high impact gene variant is present for COMT, SULT1A1 or CYP17A, the following nutritional support is recommended to effectively reduce oestrogen load by supporting preferred oestrogen pathways:

- For breakdown of oestrogen to the beneficial 2-OH metabolite, supplement with a bio-available form of 3,3'-Diindolylmethane (DIM), or substantially increase intake of cruciferous vegetables (cauliflower, broccoli, cabbage, brussels sprouts).
- Include phytoestrogens in the diet for their many beneficial influences on oestrogen synthesis and metabolism. These include isoflavones and lignins. Isoflavones are found most commonly in soy products, but also include legumes, alfalfa, clover, licorice root, and kudzu root, and include genistein, daidzein, equol and puerarin. Lignins are an insoluble dietary fibre found in flaxseeds, whole grains, beans and seeds.
- Ensure adequate intake of magnesium and Vitamin E.
- Other beneficial micro and phyto-nutrients that impact oestrogen metabolism include calcium D-glucarate, curcumin, green tea polyphenols and D-limonene.

